

Lattices and Orders in Isabelle/HOL

Markus Wenzel
TU München

June 21, 2010

Abstract

We consider abstract structures of orders and lattices. Many fundamental concepts of lattice theory are developed, including dual structures, properties of bounds versus algebraic laws, lattice operations versus set-theoretic ones etc. We also give example instantiations of lattices and orders, such as direct products and function spaces. Well-known properties are demonstrated, like the Knaster-Tarski Theorem for complete lattices.

This formal theory development may serve as an example of applying Isabelle/HOL to the domain of mathematical reasoning about “axiomatic” structures. Apart from the simply-typed classical set-theory of HOL, we employ Isabelle’s system of axiomatic type classes for expressing structures and functors in a light-weight manner. Proofs are expressed in the Isar language for readable formal proof, while aiming at its “best-style” of representing formal reasoning.

Contents

| | | |
|----------|--|----------|
| 1 | Orders | 3 |
| 1.1 | Ordered structures | 3 |
| 1.2 | Duality | 3 |
| 1.3 | Transforming orders | 5 |
| 1.3.1 | Duals | 5 |
| 1.3.2 | Binary products | 5 |
| 1.3.3 | General products | 5 |
| 2 | Bounds | 6 |
| 2.1 | Infimum and supremum | 6 |
| 2.2 | Duality | 8 |
| 2.3 | Uniqueness | 8 |
| 2.4 | Related elements | 8 |
| 2.5 | General versus binary bounds | 9 |
| 2.6 | Connecting general bounds | 9 |

| | | |
|----------|---|-----------|
| 3 | Lattices | 9 |
| 3.1 | Lattice operations | 9 |
| 3.2 | Duality | 11 |
| 3.3 | Algebraic properties | 11 |
| 3.4 | Order versus algebraic structure | 12 |
| 3.5 | Example instances | 12 |
| 3.5.1 | Linear orders | 12 |
| 3.5.2 | Binary products | 13 |
| 3.5.3 | General products | 13 |
| 3.6 | Monotonicity and semi-morphisms | 14 |
| 4 | Complete lattices | 14 |
| 4.1 | Complete lattice operations | 14 |
| 4.2 | The Knaster-Tarski Theorem | 16 |
| 4.3 | Bottom and top elements | 16 |
| 4.4 | Duality | 16 |
| 4.5 | Complete lattices are lattices | 17 |
| 4.6 | Complete lattices and set-theory operations | 17 |

1 Orders

theory *Orders* **imports** *Main* **begin**

1.1 Ordered structures

We define several classes of ordered structures over some type $'a$ with relation $\sqsubseteq :: 'a \Rightarrow 'a \Rightarrow \text{bool}$. For a *quasi-order* that relation is required to be reflexive and transitive, for a *partial order* it also has to be anti-symmetric, while for a *linear order* all elements are required to be related (in either direction).

```
class leq =
  fixes leq :: 'a  $\Rightarrow$  'a  $\Rightarrow$  bool (infixl [= 50])
```

```
notation (xsymbols)
  leq (infixl  $\sqsubseteq$  50)
```

```
class quasi-order = leq +
  assumes leq-refl [intro?]:  $x \sqsubseteq x$ 
  assumes leq-trans [trans]:  $x \sqsubseteq y \Longrightarrow y \sqsubseteq z \Longrightarrow x \sqsubseteq z$ 
```

```
class partial-order = quasi-order +
  assumes leq-antisym [trans]:  $x \sqsubseteq y \Longrightarrow y \sqsubseteq x \Longrightarrow x = y$ 
```

```
class linear-order = partial-order +
  assumes leq-linear:  $x \sqsubseteq y \vee y \sqsubseteq x$ 
```

```
lemma linear-order-cases:
   $((x :: 'a :: \text{linear-order}) \sqsubseteq y \Longrightarrow C) \Longrightarrow (y \sqsubseteq x \Longrightarrow C) \Longrightarrow C$ 
   $\langle \text{proof} \rangle$ 
```

1.2 Duality

The *dual* of an ordered structure is an isomorphic copy of the underlying type, with the \sqsubseteq relation defined as the inverse of the original one.

```
datatype 'a dual = dual 'a
```

```
consts
  undual :: 'a dual  $\Rightarrow$  'a
```

```
primrec
  undual-dual: undual (dual  $x$ ) =  $x$ 
```

```
instantiation dual :: (leq) leq
begin
```

```
definition
  leq-dual-def:  $x' \sqsubseteq y' \equiv \text{undual } y' \sqsubseteq \text{undual } x'$ 
```

```
instance  $\langle \text{proof} \rangle$ 
```

end

lemma *undual-leq* [iff?]: $(undual\ x' \sqsubseteq undual\ y') = (y' \sqsubseteq x')$
 $\langle proof \rangle$

lemma *dual-leq* [iff?]: $(dual\ x \sqsubseteq dual\ y) = (y \sqsubseteq x)$
 $\langle proof \rangle$

Functions *dual* and *undual* are inverse to each other; this entails the following fundamental properties.

lemma *dual-undual* [simp]: $dual\ (undual\ x') = x'$
 $\langle proof \rangle$

lemma *undual-dual-id* [simp]: $undual\ o\ dual = id$
 $\langle proof \rangle$

lemma *dual-undual-id* [simp]: $dual\ o\ undual = id$
 $\langle proof \rangle$

Since *dual* (and *undual*) are both injective and surjective, the basic logical connectives (equality, quantification etc.) are transferred as follows.

lemma *undual-equality* [iff?]: $(undual\ x' = undual\ y') = (x' = y')$
 $\langle proof \rangle$

lemma *dual-equality* [iff?]: $(dual\ x = dual\ y) = (x = y)$
 $\langle proof \rangle$

lemma *dual-ball* [iff?]: $(\forall x \in A. P\ (dual\ x)) = (\forall x' \in dual\ ^\circ A. P\ x')$
 $\langle proof \rangle$

lemma *range-dual* [simp]: $dual\ ^\circ UNIV = UNIV$
 $\langle proof \rangle$

lemma *dual-all* [iff?]: $(\forall x. P\ (dual\ x)) = (\forall x'. P\ x')$
 $\langle proof \rangle$

lemma *dual-ex*: $(\exists x. P\ (dual\ x)) = (\exists x'. P\ x')$
 $\langle proof \rangle$

lemma *dual-Collect*: $\{dual\ x \mid x. P\ (dual\ x)\} = \{x'. P\ x'\}$
 $\langle proof \rangle$

1.3 Transforming orders

1.3.1 Duals

The classes of quasi, partial, and linear orders are all closed under formation of dual structures.

instance *dual* :: (*quasi-order*) *quasi-order*
 $\langle proof \rangle$

instance *dual* :: (*partial-order*) *partial-order*
 $\langle proof \rangle$

instance *dual* :: (*linear-order*) *linear-order*
 $\langle proof \rangle$

1.3.2 Binary products

The classes of quasi and partial orders are closed under binary products. Note that the direct product of linear orders need *not* be linear in general.

instantiation * :: (*leq*, *leq*) *leq*
begin

definition

leq-prod-def: $p \sqsubseteq q \equiv fst\ p \sqsubseteq fst\ q \wedge snd\ p \sqsubseteq snd\ q$

instance $\langle proof \rangle$

end

lemma *leq-prodI* [*intro?*]:

$fst\ p \sqsubseteq fst\ q \implies snd\ p \sqsubseteq snd\ q \implies p \sqsubseteq q$
 $\langle proof \rangle$

lemma *leq-prodE* [*elim?*]:

$p \sqsubseteq q \implies (fst\ p \sqsubseteq fst\ q \implies snd\ p \sqsubseteq snd\ q \implies C) \implies C$
 $\langle proof \rangle$

instance * :: (*quasi-order*, *quasi-order*) *quasi-order*
 $\langle proof \rangle$

instance * :: (*partial-order*, *partial-order*) *partial-order*
 $\langle proof \rangle$

1.3.3 General products

The classes of quasi and partial orders are closed under general products (function spaces). Note that the direct product of linear orders need *not* be linear in general.

```

instantiation fun :: (type, leq) leq
begin

definition
  leq-fun-def:  $f \sqsubseteq g \equiv \forall x. f\ x \sqsubseteq g\ x$ 

instance ⟨proof⟩

end

lemma leq-funI [intro?]:  $(\bigwedge x. f\ x \sqsubseteq g\ x) \implies f \sqsubseteq g$ 
  ⟨proof⟩

lemma leq-funD [dest?]:  $f \sqsubseteq g \implies f\ x \sqsubseteq g\ x$ 
  ⟨proof⟩

instance fun :: (type, quasi-order) quasi-order
  ⟨proof⟩

instance fun :: (type, partial-order) partial-order
  ⟨proof⟩

end

```

2 Bounds

```

theory Bounds imports Orders begin

```

```

hide-const (open) inf sup

```

2.1 Infimum and supremum

Given a partial order, we define infimum (greatest lower bound) and supremum (least upper bound) wrt. \sqsubseteq for two and for any number of elements.

definition

```

is-inf :: 'a::partial-order  $\Rightarrow$  'a  $\Rightarrow$  'a  $\Rightarrow$  bool where
is-inf x y inf = (inf  $\sqsubseteq$  x  $\wedge$  inf  $\sqsubseteq$  y  $\wedge$  ( $\forall z. z \sqsubseteq x \wedge z \sqsubseteq y \longrightarrow z \sqsubseteq$  inf))

```

definition

```

is-sup :: 'a::partial-order  $\Rightarrow$  'a  $\Rightarrow$  'a  $\Rightarrow$  bool where
is-sup x y sup = (x  $\sqsubseteq$  sup  $\wedge$  y  $\sqsubseteq$  sup  $\wedge$  ( $\forall z. x \sqsubseteq z \wedge y \sqsubseteq z \longrightarrow sup \sqsubseteq$  z))

```

definition

```

is-Inf :: 'a::partial-order set  $\Rightarrow$  'a  $\Rightarrow$  bool where
is-Inf A inf = (( $\forall x \in A. inf \sqsubseteq x$ )  $\wedge$  ( $\forall z. (\forall x \in A. z \sqsubseteq x) \longrightarrow z \sqsubseteq$  inf))

```

definition

$is-Sup :: 'a::partial-order set \Rightarrow 'a \Rightarrow bool$ **where**
 $is-Sup A sup = ((\forall x \in A. x \sqsubseteq sup) \wedge (\forall z. (\forall x \in A. x \sqsubseteq z) \longrightarrow sup \sqsubseteq z))$

These definitions entail the following basic properties of boundary elements.

lemma *is-infI* [intro?]: $inf \sqsubseteq x \Longrightarrow inf \sqsubseteq y \Longrightarrow$
 $(\bigwedge z. z \sqsubseteq x \Longrightarrow z \sqsubseteq y \Longrightarrow z \sqsubseteq inf) \Longrightarrow is-inf\ x\ y\ inf$
 $\langle proof \rangle$

lemma *is-inf-greatest* [elim?]:
 $is-inf\ x\ y\ inf \Longrightarrow z \sqsubseteq x \Longrightarrow z \sqsubseteq y \Longrightarrow z \sqsubseteq inf$
 $\langle proof \rangle$

lemma *is-inf-lower* [elim?]:
 $is-inf\ x\ y\ inf \Longrightarrow (inf \sqsubseteq x \Longrightarrow inf \sqsubseteq y \Longrightarrow C) \Longrightarrow C$
 $\langle proof \rangle$

lemma *is-supI* [intro?]: $x \sqsubseteq sup \Longrightarrow y \sqsubseteq sup \Longrightarrow$
 $(\bigwedge z. x \sqsubseteq z \Longrightarrow y \sqsubseteq z \Longrightarrow sup \sqsubseteq z) \Longrightarrow is-sup\ x\ y\ sup$
 $\langle proof \rangle$

lemma *is-sup-least* [elim?]:
 $is-sup\ x\ y\ sup \Longrightarrow x \sqsubseteq z \Longrightarrow y \sqsubseteq z \Longrightarrow sup \sqsubseteq z$
 $\langle proof \rangle$

lemma *is-sup-upper* [elim?]:
 $is-sup\ x\ y\ sup \Longrightarrow (x \sqsubseteq sup \Longrightarrow y \sqsubseteq sup \Longrightarrow C) \Longrightarrow C$
 $\langle proof \rangle$

lemma *is-InfI* [intro?]: $(\bigwedge x. x \in A \Longrightarrow inf \sqsubseteq x) \Longrightarrow$
 $(\bigwedge z. (\forall x \in A. z \sqsubseteq x) \Longrightarrow z \sqsubseteq inf) \Longrightarrow is-Inf\ A\ inf$
 $\langle proof \rangle$

lemma *is-Inf-greatest* [elim?]:
 $is-Inf\ A\ inf \Longrightarrow (\bigwedge x. x \in A \Longrightarrow z \sqsubseteq x) \Longrightarrow z \sqsubseteq inf$
 $\langle proof \rangle$

lemma *is-Inf-lower* [dest?]:
 $is-Inf\ A\ inf \Longrightarrow x \in A \Longrightarrow inf \sqsubseteq x$
 $\langle proof \rangle$

lemma *is-SupI* [intro?]: $(\bigwedge x. x \in A \Longrightarrow x \sqsubseteq sup) \Longrightarrow$
 $(\bigwedge z. (\forall x \in A. x \sqsubseteq z) \Longrightarrow sup \sqsubseteq z) \Longrightarrow is-Sup\ A\ sup$
 $\langle proof \rangle$

lemma *is-Sup-least* [elim?]:
 $is-Sup\ A\ sup \Longrightarrow (\bigwedge x. x \in A \Longrightarrow x \sqsubseteq z) \Longrightarrow sup \sqsubseteq z$

$\langle proof \rangle$

lemma *is-Sup-upper* [*dest?*]:

$$is-Sup\ A\ sup \implies x \in A \implies x \sqsubseteq sup$$

$\langle proof \rangle$

2.2 Duality

Infimum and supremum are dual to each other.

theorem *dual-inf* [*iff?*]:

$$is-inf\ (dual\ x)\ (dual\ y)\ (dual\ sup) = is-sup\ x\ y\ sup$$

$\langle proof \rangle$

theorem *dual-sup* [*iff?*]:

$$is-sup\ (dual\ x)\ (dual\ y)\ (dual\ inf) = is-inf\ x\ y\ inf$$

$\langle proof \rangle$

theorem *dual-Inf* [*iff?*]:

$$is-Inf\ (dual\ 'A)\ (dual\ sup) = is-Sup\ A\ sup$$

$\langle proof \rangle$

theorem *dual-Sup* [*iff?*]:

$$is-Sup\ (dual\ 'A)\ (dual\ inf) = is-Inf\ A\ inf$$

$\langle proof \rangle$

2.3 Uniqueness

Infima and suprema on partial orders are unique; this is mainly due to anti-symmetry of the underlying relation.

theorem *is-inf-uniq*: $is-inf\ x\ y\ inf \implies is-inf\ x\ y\ inf' \implies inf = inf'$

$\langle proof \rangle$

theorem *is-sup-uniq*: $is-sup\ x\ y\ sup \implies is-sup\ x\ y\ sup' \implies sup = sup'$

$\langle proof \rangle$

theorem *is-Inf-uniq*: $is-Inf\ A\ inf \implies is-Inf\ A\ inf' \implies inf = inf'$

$\langle proof \rangle$

theorem *is-Sup-uniq*: $is-Sup\ A\ sup \implies is-Sup\ A\ sup' \implies sup = sup'$

$\langle proof \rangle$

2.4 Related elements

The binary bound of related elements is either one of the argument.

theorem *is-inf-related* [*elim?*]: $x \sqsubseteq y \implies is-inf\ x\ y\ x$

$\langle proof \rangle$

theorem *is-sup-related* [*elim?*]: $x \sqsubseteq y \implies \text{is-sup } x \ y$
<proof>

2.5 General versus binary bounds

General bounds of two-element sets coincide with binary bounds.

theorem *is-Inf-binary*: $\text{is-Inf } \{x, y\} \ \text{inf} = \text{is-inf } x \ y \ \text{inf}$
<proof>

theorem *is-Sup-binary*: $\text{is-Sup } \{x, y\} \ \text{sup} = \text{is-sup } x \ y \ \text{sup}$
<proof>

2.6 Connecting general bounds

Either kind of general bounds is sufficient to express the other. The least upper bound (supremum) is the same as the the greatest lower bound of the set of all upper bounds; the dual statements holds as well; the dual statement holds as well.

theorem *Inf-Sup*: $\text{is-Inf } \{b. \forall a \in A. a \sqsubseteq b\} \ \text{sup} \implies \text{is-Sup } A \ \text{sup}$
<proof>

theorem *Sup-Inf*: $\text{is-Sup } \{b. \forall a \in A. b \sqsubseteq a\} \ \text{inf} \implies \text{is-Inf } A \ \text{inf}$
<proof>

end

3 Lattices

theory *Lattice* **imports** *Bounds* **begin**

3.1 Lattice operations

A *lattice* is a partial order with infimum and supremum of any two elements (thus any *finite* number of elements have bounds as well).

class *lattice* =
assumes *ex-inf*: $\exists \text{inf}. \text{is-inf } x \ y \ \text{inf}$
assumes *ex-sup*: $\exists \text{sup}. \text{is-sup } x \ y \ \text{sup}$

The \sqcap (meet) and \sqcup (join) operations select such infimum and supremum elements.

definition
 $\text{meet} :: 'a::\text{lattice} \Rightarrow 'a \Rightarrow 'a$ (**infixl** $\&\&$ 70) **where**
 $x \ \&\& \ y = (\text{THE } \text{inf}. \text{is-inf } x \ y \ \text{inf})$

definition
 $\text{join} :: 'a::\text{lattice} \Rightarrow 'a \Rightarrow 'a$ (**infixl** $\|\$ 65) **where**

$$x \parallel y = (\text{THE } \text{sup. is-sup } x \ y \ \text{sup})$$

notation (*xsymbols*)
meet (**infixl** \sqcap 70) **and**
join (**infixl** \sqcup 65)

Due to unique existence of bounds, the lattice operations may be exhibited as follows.

lemma *meet-equality* [*elim?*]: $\text{is-inf } x \ y \ \text{inf} \implies x \sqcap y = \text{inf}$
 $\langle \text{proof} \rangle$

lemma *meetI* [*intro?*]:
 $\text{inf} \sqsubseteq x \implies \text{inf} \sqsubseteq y \implies (\bigwedge z. z \sqsubseteq x \implies z \sqsubseteq y \implies z \sqsubseteq \text{inf}) \implies x \sqcap y = \text{inf}$
 $\langle \text{proof} \rangle$

lemma *join-equality* [*elim?*]: $\text{is-sup } x \ y \ \text{sup} \implies x \sqcup y = \text{sup}$
 $\langle \text{proof} \rangle$

lemma *joinI* [*intro?*]: $x \sqsubseteq \text{sup} \implies y \sqsubseteq \text{sup} \implies$
 $(\bigwedge z. x \sqsubseteq z \implies y \sqsubseteq z \implies \text{sup} \sqsubseteq z) \implies x \sqcup y = \text{sup}$
 $\langle \text{proof} \rangle$

The \sqcap and \sqcup operations indeed determine bounds on a lattice structure.

lemma *is-inf-meet* [*intro?*]: $\text{is-inf } x \ y \ (x \sqcap y)$
 $\langle \text{proof} \rangle$

lemma *meet-greatest* [*intro?*]: $z \sqsubseteq x \implies z \sqsubseteq y \implies z \sqsubseteq x \sqcap y$
 $\langle \text{proof} \rangle$

lemma *meet-lower1* [*intro?*]: $x \sqcap y \sqsubseteq x$
 $\langle \text{proof} \rangle$

lemma *meet-lower2* [*intro?*]: $x \sqcap y \sqsubseteq y$
 $\langle \text{proof} \rangle$

lemma *is-sup-join* [*intro?*]: $\text{is-sup } x \ y \ (x \sqcup y)$
 $\langle \text{proof} \rangle$

lemma *join-least* [*intro?*]: $x \sqsubseteq z \implies y \sqsubseteq z \implies x \sqcup y \sqsubseteq z$
 $\langle \text{proof} \rangle$

lemma *join-upper1* [*intro?*]: $x \sqsubseteq x \sqcup y$
 $\langle \text{proof} \rangle$

lemma *join-upper2* [*intro?*]: $y \sqsubseteq x \sqcup y$
 $\langle \text{proof} \rangle$

3.2 Duality

The class of lattices is closed under formation of dual structures. This means that for any theorem of lattice theory, the dualized statement holds as well; this important fact simplifies many proofs of lattice theory.

instance *dual* :: (*lattice*) *lattice*
 ⟨*proof*⟩

Apparently, the \sqcap and \sqcup operations are dual to each other.

theorem *dual-meet* [*intro?*]: $\text{dual } (x \sqcap y) = \text{dual } x \sqcup \text{dual } y$
 ⟨*proof*⟩

theorem *dual-join* [*intro?*]: $\text{dual } (x \sqcup y) = \text{dual } x \sqcap \text{dual } y$
 ⟨*proof*⟩

3.3 Algebraic properties

The \sqcap and \sqcup operations have the following characteristic algebraic properties: associative (A), commutative (C), and absorptive (AB).

theorem *meet-assoc*: $(x \sqcap y) \sqcap z = x \sqcap (y \sqcap z)$
 ⟨*proof*⟩

theorem *join-assoc*: $(x \sqcup y) \sqcup z = x \sqcup (y \sqcup z)$
 ⟨*proof*⟩

theorem *meet-commute*: $x \sqcap y = y \sqcap x$
 ⟨*proof*⟩

theorem *join-commute*: $x \sqcup y = y \sqcup x$
 ⟨*proof*⟩

theorem *meet-join-absorb*: $x \sqcap (x \sqcup y) = x$
 ⟨*proof*⟩

theorem *join-meet-absorb*: $x \sqcup (x \sqcap y) = x$
 ⟨*proof*⟩

Some further algebraic properties hold as well. The property idempotent (I) is a basic algebraic consequence of (AB).

theorem *meet-idem*: $x \sqcap x = x$
 ⟨*proof*⟩

theorem *join-idem*: $x \sqcup x = x$
 ⟨*proof*⟩

Meet and join are trivial for related elements.

theorem *meet-related* [*elim?*]: $x \sqsubseteq y \implies x \sqcap y = x$
 $\langle \text{proof} \rangle$

theorem *join-related* [*elim?*]: $x \sqsubseteq y \implies x \sqcup y = y$
 $\langle \text{proof} \rangle$

3.4 Order versus algebraic structure

The \sqcap and \sqcup operations are connected with the underlying \sqsubseteq relation in a canonical manner.

theorem *meet-connection*: $(x \sqsubseteq y) = (x \sqcap y = x)$
 $\langle \text{proof} \rangle$

theorem *join-connection*: $(x \sqsubseteq y) = (x \sqcup y = y)$
 $\langle \text{proof} \rangle$

The most fundamental result of the meta-theory of lattices is as follows (we do not prove it here).

Given a structure with binary operations \sqcap and \sqcup such that (A), (C), and (AB) hold (cf. §3.3). This structure represents a lattice, if the relation $x \sqsubseteq y$ is defined as $x \sqcap y = x$ (alternatively as $x \sqcup y = y$). Furthermore, infimum and supremum with respect to this ordering coincide with the original \sqcap and \sqcup operations.

3.5 Example instances

3.5.1 Linear orders

Linear orders with *minimum* and *maximum* operations are a (degenerate) example of lattice structures.

definition

minimum :: 'a::linear-order \Rightarrow 'a \Rightarrow 'a **where**
minimum x y = (if x \sqsubseteq y then x else y)

definition

maximum :: 'a::linear-order \Rightarrow 'a \Rightarrow 'a **where**
maximum x y = (if x \sqsubseteq y then y else x)

lemma *is-inf-minimum*: *is-inf* x y (*minimum* x y)
 $\langle \text{proof} \rangle$

lemma *is-sup-maximum*: *is-sup* x y (*maximum* x y)
 $\langle \text{proof} \rangle$

instance *linear-order* \subseteq *lattice*
 $\langle \text{proof} \rangle$

The lattice operations on linear orders indeed coincide with *minimum* and *maximum*.

theorem *meet-minimum*: $x \sqcap y = \text{minimum } x \ y$
 $\langle \text{proof} \rangle$

theorem *meet-maximum*: $x \sqcup y = \text{maximum } x \ y$
 $\langle \text{proof} \rangle$

3.5.2 Binary products

The class of lattices is closed under direct binary products (cf. §1.3.2).

lemma *is-inf-prod*: $\text{is-inf } p \ q \ (\text{fst } p \sqcap \text{fst } q, \text{snd } p \sqcap \text{snd } q)$
 $\langle \text{proof} \rangle$

lemma *is-sup-prod*: $\text{is-sup } p \ q \ (\text{fst } p \sqcup \text{fst } q, \text{snd } p \sqcup \text{snd } q)$
 $\langle \text{proof} \rangle$

instance $*$:: (lattice, lattice) lattice
 $\langle \text{proof} \rangle$

The lattice operations on a binary product structure indeed coincide with the products of the original ones.

theorem *meet-prod*: $p \sqcap q = (\text{fst } p \sqcap \text{fst } q, \text{snd } p \sqcap \text{snd } q)$
 $\langle \text{proof} \rangle$

theorem *join-prod*: $p \sqcup q = (\text{fst } p \sqcup \text{fst } q, \text{snd } p \sqcup \text{snd } q)$
 $\langle \text{proof} \rangle$

3.5.3 General products

The class of lattices is closed under general products (function spaces) as well (cf. §1.3.3).

lemma *is-inf-fun*: $\text{is-inf } f \ g \ (\lambda x. \text{fst } f \ x \sqcap \text{fst } g \ x)$
 $\langle \text{proof} \rangle$

lemma *is-sup-fun*: $\text{is-sup } f \ g \ (\lambda x. \text{fst } f \ x \sqcup \text{fst } g \ x)$
 $\langle \text{proof} \rangle$

instance *fun* :: (type, lattice) lattice
 $\langle \text{proof} \rangle$

The lattice operations on a general product structure (function space) indeed emerge by point-wise lifting of the original ones.

theorem *meet-fun*: $f \sqcap g = (\lambda x. \text{fst } f \ x \sqcap \text{fst } g \ x)$
 $\langle \text{proof} \rangle$

theorem *join-fun*: $f \sqcup g = (\lambda x. f x \sqcup g x)$
 $\langle proof \rangle$

3.6 Monotonicity and semi-morphisms

The lattice operations are monotone in both argument positions. In fact, monotonicity of the second position is trivial due to commutativity.

theorem *meet-mono*: $x \sqsubseteq z \implies y \sqsubseteq w \implies x \sqcap y \sqsubseteq z \sqcap w$
 $\langle proof \rangle$

theorem *join-mono*: $x \sqsubseteq z \implies y \sqsubseteq w \implies x \sqcup y \sqsubseteq z \sqcup w$
 $\langle proof \rangle$

A semi-morphisms is a function f that preserves the lattice operations in the following manner: $f (x \sqcap y) \sqsubseteq f x \sqcap f y$ and $f x \sqcup f y \sqsubseteq f (x \sqcup y)$, respectively. Any of these properties is equivalent with monotonicity.

theorem *meet-semimorph*:
 $(\bigwedge x y. f (x \sqcap y) \sqsubseteq f x \sqcap f y) \equiv (\bigwedge x y. x \sqsubseteq y \implies f x \sqsubseteq f y)$
 $\langle proof \rangle$

lemma *join-semimorph*:
 $(\bigwedge x y. f x \sqcup f y \sqsubseteq f (x \sqcup y)) \equiv (\bigwedge x y. x \sqsubseteq y \implies f x \sqsubseteq f y)$
 $\langle proof \rangle$

end

4 Complete lattices

theory *CompleteLattice* **imports** *Lattice* **begin**

4.1 Complete lattice operations

A *complete lattice* is a partial order with general (infinitary) infimum of any set of elements. General supremum exists as well, as a consequence of the connection of infinitary bounds (see §2.6).

class *complete-lattice* =
assumes *ex-Inf*: $\exists inf. is-Inf A inf$

theorem *ex-Sup*: $\exists sup::'a::complete-lattice. is-Sup A sup$
 $\langle proof \rangle$

The general \sqcap (meet) and \sqcup (join) operations select such infimum and supremum elements.

definition

Meet :: 'a::complete-lattice set \Rightarrow 'a **where**
Meet A = (THE inf. is-Inf A inf)

definition

Join :: 'a::complete-lattice set \Rightarrow 'a **where**
Join A = (THE sup. is-Sup A sup)

notation (xsymbols)

Meet (\sqcap - [90] 90) **and**
Join (\sqcup - [90] 90)

Due to unique existence of bounds, the complete lattice operations may be exhibited as follows.

lemma *Meet-equality* [elim?]: is-Inf A inf $\Longrightarrow \sqcap A = \text{inf}$
 <proof>

lemma *MeetI* [intro?]:
 ($\bigwedge a. a \in A \Longrightarrow \text{inf} \sqsubseteq a$) \Longrightarrow
 ($\bigwedge b. \forall a \in A. b \sqsubseteq a \Longrightarrow b \sqsubseteq \text{inf}$) \Longrightarrow
 $\sqcap A = \text{inf}$
 <proof>

lemma *Join-equality* [elim?]: is-Sup A sup $\Longrightarrow \sqcup A = \text{sup}$
 <proof>

lemma *JoinI* [intro?]:
 ($\bigwedge a. a \in A \Longrightarrow a \sqsubseteq \text{sup}$) \Longrightarrow
 ($\bigwedge b. \forall a \in A. a \sqsubseteq b \Longrightarrow \text{sup} \sqsubseteq b$) \Longrightarrow
 $\sqcup A = \text{sup}$
 <proof>

The \sqcap and \sqcup operations indeed determine bounds on a complete lattice structure.

lemma *is-Inf-Meet* [intro?]: is-Inf A ($\sqcap A$)
 <proof>

lemma *Meet-greatest* [intro?]: ($\bigwedge a. a \in A \Longrightarrow x \sqsubseteq a$) $\Longrightarrow x \sqsubseteq \sqcap A$
 <proof>

lemma *Meet-lower* [intro?]: $a \in A \Longrightarrow \sqcap A \sqsubseteq a$
 <proof>

lemma *is-Sup-Join* [intro?]: is-Sup A ($\sqcup A$)
 <proof>

lemma *Join-least* [intro?]: ($\bigwedge a. a \in A \Longrightarrow a \sqsubseteq x$) $\Longrightarrow \sqcup A \sqsubseteq x$
 <proof>

lemma *Join-lower* [intro?]: $a \in A \Longrightarrow a \sqsubseteq \sqcup A$

$\langle proof \rangle$

4.2 The Knaster-Tarski Theorem

The Knaster-Tarski Theorem (in its simplest formulation) states that any monotone function on a complete lattice has a least fixed-point (see [2, pages 93–94] for example). This is a consequence of the basic boundary properties of the complete lattice operations.

theorem *Knaster-Tarski*:

assumes *mono*: $\bigwedge x y. x \sqsubseteq y \implies f x \sqsubseteq f y$

obtains $a :: 'a::complete-lattice$ **where**

$f a = a$ **and** $\bigwedge a'. f a' = a' \implies a \sqsubseteq a'$

$\langle proof \rangle$

theorem *Knaster-Tarski-dual*:

assumes *mono*: $\bigwedge x y. x \sqsubseteq y \implies f x \sqsubseteq f y$

obtains $a :: 'a::complete-lattice$ **where**

$f a = a$ **and** $\bigwedge a'. f a' = a' \implies a' \sqsubseteq a$

$\langle proof \rangle$

4.3 Bottom and top elements

With general bounds available, complete lattices also have least and greatest elements.

definition

bottom :: $'a::complete-lattice$ (\perp) **where**

$\perp = \bigcap UNIV$

definition

top :: $'a::complete-lattice$ (\top) **where**

$\top = \bigcup UNIV$

lemma *bottom-least* $[intro?]$: $\perp \sqsubseteq x$

$\langle proof \rangle$

lemma *bottomI* $[intro?]$: $(\bigwedge a. x \sqsubseteq a) \implies \perp = x$

$\langle proof \rangle$

lemma *top-greatest* $[intro?]$: $x \sqsubseteq \top$

$\langle proof \rangle$

lemma *topI* $[intro?]$: $(\bigwedge a. a \sqsubseteq x) \implies \top = x$

$\langle proof \rangle$

4.4 Duality

The class of complete lattices is closed under formation of dual structures.

instance *dual* :: (*complete-lattice*) *complete-lattice*
 ⟨*proof*⟩

Apparently, the \sqcap and \sqcup operations are dual to each other.

theorem *dual-Meet* [*intro?*]: *dual* ($\sqcap A$) = \sqcup (*dual* ‘ *A*)
 ⟨*proof*⟩

theorem *dual-Join* [*intro?*]: *dual* ($\sqcup A$) = \sqcap (*dual* ‘ *A*)
 ⟨*proof*⟩

Likewise are \perp and \top duals of each other.

theorem *dual-bottom* [*intro?*]: *dual* \perp = \top
 ⟨*proof*⟩

theorem *dual-top* [*intro?*]: *dual* \top = \perp
 ⟨*proof*⟩

4.5 Complete lattices are lattices

Complete lattices (with general bounds available) are indeed plain lattices as well. This holds due to the connection of general versus binary bounds that has been formally established in §2.5.

lemma *is-inf-binary*: *is-inf* *x y* ($\sqcap \{x, y\}$)
 ⟨*proof*⟩

lemma *is-sup-binary*: *is-sup* *x y* ($\sqcup \{x, y\}$)
 ⟨*proof*⟩

instance *complete-lattice* \subseteq *lattice*
 ⟨*proof*⟩

theorem *meet-binary*: *x* \sqcap *y* = $\sqcap \{x, y\}$
 ⟨*proof*⟩

theorem *join-binary*: *x* \sqcup *y* = $\sqcup \{x, y\}$
 ⟨*proof*⟩

4.6 Complete lattices and set-theory operations

The complete lattice operations are (anti) monotone wrt. set inclusion.

theorem *Meet-subset-antimono*: $A \subseteq B \implies \sqcap B \sqsubseteq \sqcap A$
 ⟨*proof*⟩

theorem *Join-subset-mono*: $A \subseteq B \implies \sqcup A \sqsubseteq \sqcup B$
 ⟨*proof*⟩

Bounds over unions of sets may be obtained separately.

theorem *Meet-Un*: $\prod (A \cup B) = \prod A \sqcap \prod B$
 $\langle proof \rangle$

theorem *Join-Un*: $\sqcup (A \cup B) = \sqcup A \sqcup \sqcup B$
 $\langle proof \rangle$

Bounds over singleton sets are trivial.

theorem *Meet-singleton*: $\prod \{x\} = x$
 $\langle proof \rangle$

theorem *Join-singleton*: $\sqcup \{x\} = x$
 $\langle proof \rangle$

Bounds over the empty and universal set correspond to each other.

theorem *Meet-empty*: $\prod \{\} = \sqcup UNIV$
 $\langle proof \rangle$

theorem *Join-empty*: $\sqcup \{\} = \prod UNIV$
 $\langle proof \rangle$

end

References

- [1] G. Bauer and M. Wenzel. Computer-assisted mathematics at work — the Hahn-Banach theorem in Isabelle/Isar. In T. Coquand, P. Dybjer, B. Nordström, and J. Smith, editors, *Types for Proofs and Programs: TYPES'99*, LNCS, 2000.
- [2] B. A. Davey and H. A. Priestley. *Introduction to Lattices and Order*. Cambridge University Press, 1990.
- [3] M. Wenzel. Isar — a generic interpretative approach to readable formal proof documents. In Y. Bertot, G. Dowek, A. Hirschowitz, C. Paulin, and L. Thery, editors, *Theorem Proving in Higher Order Logics: TPHOLs '99*, volume 1690 of *LNCS*, 1999.
- [4] M. Wenzel. *The Isabelle/Isar Reference Manual*, 2000. <http://isabelle.in.tum.de/doc/isar-ref.pdf>.
- [5] M. Wenzel. *Using Axiomatic Type Classes in Isabelle*, 2000. <http://isabelle.in.tum.de/doc/axclass.pdf>.